

Lorentz-violation and cosmological perturbations: a toy brane-world model

M.V. Libanov[†] and V.A. Rubakov[†]

[†] Institute for Nuclear Research of the Russian Academy of Sciences,
60th October Anniversary prospect 7a, Moscow 117312, Russia

E-mail: rubakov@ms2.inr.ac.ru, ml@ms2.inr.ac.ru

Abstract. We study possible effects of Lorentz-violation on the generation of cosmological perturbations at inflation by introducing a simple inflating five-dimensional brane-world setup with violation of four-dimensional Lorentz-invariance at an energy scale k . We consider massless scalar field, meant to mimic perturbations of inflaton and/or gravitational field, in this background. At three-momenta below k , there exists a zero mode localized on the brane, whose behaviour coincides with that in four-dimensional theory. On the contrary, at three-momenta above k , the localized mode is absent and physics is entirely five-dimensional. As three-momenta get redshifted, more modes get localized on the brane, the phenomenon analogous to “mode generation”. We find that for $k \gg H$, where H is the inflationary Hubble scale, the spectrum of perturbations coincides with that in four-dimensional theory. For $k < H$ and time-dependent bulk parameters, the spectrum deviates, possibly strongly, from the flat spectrum even for pure de Sitter inflation.

PACS numbers: 98.80.-k, 04.50.+h,

1. Introduction and summary

The standard inflationary mechanism of the generation of cosmological perturbations assumes Lorentz-invariance. This assumption is worth questioning [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22] especially in view of the fact that relevant perturbations initially have wavelengths much shorter than the Planck length (this observation is often referred to as “trans-Planckian problem”). Furthermore, there is no guarantee that Lorentz-invariance holds even towards the end of inflation. Thus, it is certainly of interest to understand the effects that Lorentz-violation may have on the cosmological perturbations.

Several ways to address this issue have been proposed. One is to assume Lorentz-violating dispersion relation at short enough wavelengths [1, 2, 3, 4, 5, 6, 7]. If at asymptotically high momenta the frequencies exceed the inflationary Hubble parameter, the adiabatic vacuum is a natural choice of the initial state, and then the primordial spectrum is still flat for pure de Sitter inflation [1, 2, 3, 4, 5], while the predictions for the tilt may be different, as compared to the standard mechanism, if the expansion is not exactly de Sitter. The opposite case of low frequencies at high momenta [1, 2] is hard to treat: perturbations are not initially in the adiabatic regime, so there is no preferred initial state (an exception here is the bouncing Universe model of reference [6]).

Another approach is to make ad hoc assumptions on the initial state of perturbations [8, 9, 10, 11]. The problem with this approach is that any choice other than the adiabatic vacuum is hard to justify, whereas different choices lead to quite different spectra of perturbations [10, 11].

One more possibility has to do with “mode generation”: there are models where the number of modes increases in time, as the Universe expands [12, 13, 14, 15, 16]; this happens, e.g., if the physical momentum is effectively bounded from above. This option is in a sense similar to the previous one: the result for the spectrum of perturbations strongly depends on the choice of a state in which a newborn mode emerges. If this is the adiabatic vacuum, the outcome is similar to the standard scenario, at least for pure de Sitter inflation [13, 14, 16].

The emerging picture is that in the least exotic scenarios, the spectrum of perturbations is flat for pure de Sitter inflation, while predictions for the tilt in non-de Sitter cases are generally different from those of the standard mechanism. If the energy scale k associated with Lorentz-violation is much higher than the Hubble parameter H at inflation, the deviations of the spectrum from the predictions of the standard mechanism are suppressed by powers of H/k , as argued on general grounds, e.g., in references [4, 17, 18]. On the other hand, dramatic assumptions, like, e.g., the Corley–Jacobson dispersion relation [23] with frequencies tending to zero at high momenta, lead to dramatic consequences: the spectrum may be quite different from the flat spectrum even for pure de Sitter inflation [1, 2, 7]. Also, for $k \lesssim H$ the properties of the primordial perturbations may deviate substantially from the standard predictions even in some less exotic models, like, e.g., the model of reference [19] based

on noncommutative field theory.

Violation of four-dimensional Lorentz-invariance is fairly natural in brane-world models [24, 25, 26, 27, 28]. It may occur in the bulk, especially if there is bulk matter (a particularly well-understood solution of this sort is a brane moving in Schwarzschild–anti-de Sitter bulk [29, 30]) or Lorentz-violating bulk condensate [31]. Since the wave functions of brane matter may extend to the bulk, the dispersion relations on the brane are then naturally Lorentz-violating [27]. Furthermore, some brane-world geometries have the property of “mode generation” in the following sense: for low three-momenta, there exist modes localized (or quasi-localized) on the brane, while for high three-momenta, localized and even quasi-localized modes are absent [28]. Thus, as the Universe expands and three-momenta get redshifted, more and more brane modes emerge; this may be viewed as the brane-world version of mode generation. Yet the initial state is well defined, provided that modes with high three-momenta are adiabatic in the bulk.

In this paper we consider an inflating version of the setup of reference [28], and study massless scalar field, meant to mimic perturbations of the inflaton and/or gravitational field, in this background. This is nothing more than a toy model, as we specify neither the mechanism giving rise to Lorentz-violation, nor the mechanism driving inflation. In fact, our background may be totally unrealistic, as it may require matter with unphysical properties either in the bulk or on the brane, or both. Also, since we do not have a mechanism generating our background, we cannot quantitatively address the issue of back reaction [20, 4, 21, 22] of the perturbations on background geometry (however, we will argue on qualitative grounds that this back reaction is small). On the other hand, the advantage of our approach is that the properties of the massless scalar field are well understood in spite of mode generation on the brane and other unusual features.

To a certain extent we confirm the picture emerging from the previous studies. In the simplest version of our setup, with time-independent bulk parameters, the spectrum of perturbations generated due to inflation is flat for pure de Sitter expansion, while its amplitude and, in the case of non-de Sitter expansion, the tilt are generally different from those predicted by the standard mechanism. Furthermore, for $k \gg H$, where k and H are still the scales associated with Lorentz-violation and expansion rate at inflation, respectively, even the latter deviations are small.

Things are more interesting if one makes a relatively mild assumption that the bulk parameters depend on time, so that k is smaller than H towards the end of inflation. As an example, we consider the case in which k slowly decreases in time, from $k \gg H$ at the early stage of inflation to $k < H$ towards its end. Then the primordial spectrum is flat at low momenta, gets strongly tilted at intermediate momenta, and flattens out again at high momenta, as sketched in figure 1. The position of the tilted part, as well as its width, depend on the parameters of the model; this part may well extend over several (but not necessarily many) decimal places, and the amplitudes at low and high momenta may differ also by several orders of magnitude. To avoid confusion, the primordial spectrum is cut off in the ultraviolet in the usual way, the cutoff being at

physical momenta of order H at the end of inflation.

The opposite situation, shown in figure 2, occurs if k increases in time from $k < H$ to $k \gg H$.

A lesson we learn from our toy model is twofold. On the one hand, we confirm that it is quite unlikely that in a well defined setup with Lorentz-violation well above the inflationary Hubble scale ($k \gg H$ in our notations), the standard predictions may be strongly modified. In our model, physics at three-momenta above k is entirely multi-dimensional, and yet for $k \gg H$ this has basically no effect on primordial perturbations.




Figure 1. Power of perturbations, as function of momentum p , generated in a model with k decreasing in time. $p_\times$ is a free parameter and $\epsilon \ll 1$ is another free parameter of the model. The exponent α is model dependent and varies in the range $0 < \alpha < 3/2$. See section 5.1 for explanation and qualifications.




Figure 2. Same as in figure 1 but with k increasing in time.

On the other hand, if Lorentz-violation is characterized by low energy scale $k \lesssim H$, the spectrum of perturbations strongly depends on details of the cosmological evolution at inflation, and may well be grossly different from the standard approximately flat spectrum.

The spectrum sketched in figure 1 is interesting. If our scalar field is interpreted as mimicking tensor perturbations, this spectrum corresponds to greater (maybe much greater) primordial amplitudes of gravity waves at shorter wavelengths[‡]. If so, a possible non-observation of the effects of tensor perturbations on CMB would not preclude the detection of primordial gravity waves at shorter wavelengths by techniques ranging from timing of pulsars to ground-based interferometers. Moreover, since the latter techniques cover a wide range of wavelengths, one might be able, with luck, to uncover an interesting pattern of relic stochastic gravity waves in our Universe.

Likewise, the spectrum shown in figure 2 would correspond to relatively high power at long wavelengths, possibly measurable through CMB, with deviations from flatness potentially detectable by experiments searching for gravity waves at shorter wavelengths.

We conclude this section by mentioning that even more complicated spectra are obtained if one allows for time-dependence not only of k , but also of other bulk parameters. We will comment on this point in section 5.

2. Setup

We consider (4+1)-dimensional model and denote the coordinates as (η, x^i, y) , $i = 1, 2, 3$. We take the five-dimensional metric in the following form (signature $+, -, -, -, -$),

$$ds^2 = a^2(\eta)[\alpha^2(y)d\eta^2 - \beta^2(y)d\mathbf{x}^2] - \alpha^2(y)dy^2 \quad (1)$$

where we made a convenient choice of the coordinate y . There is a single brane at

$$y = y_b = 0$$

As usual, the warp factors $\alpha(y)$ and $\beta(y)$ are continuous across the brane, while their derivatives $\partial_y \alpha$ and $\partial_y \beta$ are not. For equal warp factors, $\alpha(y) = \beta(y)$, and static geometry $a(\eta) = \text{const}$, equation (1) has a general form of warped metric invariant under four-dimensional Lorentz group. For $\alpha(y) \neq \beta(y)$, and still $a(\eta) = \text{const}$, the coordinates can be chosen in such a way that the induced metric on the brane is still four-dimensionally Lorentz-invariant, while the full five-dimensional metric is not. This is the case studied, e.g., in references [27, 28]. In what follows we choose coordinates so that the brane is Lorentz-invariant for $a = \text{const}$. To specify our setup further, we assume that both $\alpha(y)$ and $\beta(y)$ are Z_2 -symmetric across the brane and decay away from the brane,

$$\alpha(y), \beta(y) \rightarrow 0, \quad \text{as } y \rightarrow \infty$$

and that their ratio tends to a small constant,

$$\frac{\alpha(y)}{\beta(y)} \rightarrow \epsilon, \quad \text{as } y \rightarrow \infty \quad (2)$$

[‡] As usual, the gravity waves decay as a^{-1} after re-entering the horizon.

$$\epsilon \ll 1$$

The case $\epsilon = 0$ and $a(\eta) = \text{const}$ was discussed in reference [28].

We are interested in inflating setup. For de Sitter inflation, which is the main focus in this paper, one has

$$a(\eta) = -\frac{1}{H\eta} \quad (3)$$

If inflation is not exactly de Sitter, the Hubble parameter H slowly varies in time. Later on we will also consider generalizations of the setup (1) in which α and/or β slowly vary in time as well.

Throughout this paper we concentrate on the properties of five-dimensional massless minimal scalar field ϕ in the background (1). Its action is

$$S_\phi = \frac{1}{2} \int d^5X \sqrt{g} g^{AB} \partial_A \phi \partial_B \phi$$

Upon introducing another field χ via

$$\phi = \frac{1}{a\beta^{3/2}} \chi \quad (4)$$

the action becomes

$$S_\chi = \frac{1}{2} \int d\eta d^3x dy \left\{ \dot{\chi}^2 + \frac{\ddot{a}}{a} \chi^2 - \frac{\alpha^2(y)}{\beta^2(y)} (\partial_{\mathbf{x}} \chi)^2 - a^2(\eta) \left[(\chi')^2 + \left(\frac{3}{2} \frac{\beta''}{\beta} + \frac{3}{4} \frac{\beta'^2}{\beta^2} \right) \chi^2 \right] \right\}$$

where dot and prime denote ∂_η and ∂_y , respectively. The field equation in terms of three-dimensional Fourier harmonics is

$$-\ddot{\chi} + \frac{\ddot{a}}{a} \chi - U(y) p^2 \chi - a^2(\eta) [-\chi'' + V(y) \chi] = 0 \quad (5)$$

where p is conformal three-momentum, and the “potentials” are

$$U(y) = \frac{\alpha^2(y)}{\beta^2(y)}$$

and

$$V(y) = \frac{3}{2} \frac{\beta''}{\beta} + \frac{3}{4} \frac{\beta'^2}{\beta^2}$$

In what follows we assume that $V(y)$ tends to a constant as $|y| \rightarrow \infty$, which, for future convenience, we denote as

$$V(|y| \rightarrow \infty) = \frac{9}{4} k^2$$

$U(y)$ is a monotonically decreasing function whose asymptotics, according to equation (2), is

$$U(|y| \rightarrow \infty) = \epsilon^2$$

One more assumption is that the width y_0 of the “potential” $U(y)$, i.e., the size of the region where $U(y)$ substantially exceeds ϵ^2 , is sufficiently small; roughly speaking,

$$y_0 < k^{-1} \quad (6)$$

The reason for this assumption will become clear later[§]. Finally, we will assume that k and ϵ are the only relevant parameters entering $U(y)$ and $V(y)$, the width y_0 being roughly of order k^{-1} or maybe substantially smaller.

We have to specify the boundary condition on the brane. We impose Z_2 symmetry across the brane, consider the theory on half-space $y \geq 0$, and, guided by the properties of gravitational perturbations in the Randall–Sundrum model (see, e.g., [32] and references therein), impose the condition

$$\phi'(y=0) = 0$$

i.e.,

$$\left(\chi' - \frac{3}{2} \frac{\beta'}{\beta} \chi \right)_{y=0} = 0 \quad (7)$$

This is in accord with the discontinuity of $\beta'(y)$ at $y = 0$, which results in the δ -function behaviour of $V(y)$,

$$V(y) \approx 3 \frac{\beta'}{\beta} \delta(y) \quad \text{at } y \approx 0 \quad (8)$$

Hereafter we assume that $\beta' < 0$ for all y , so that this contribution to the potential is attractive.

Although our analysis will not depend on the precise form of the warp factors $\alpha(y)$ and $\beta(y)$, in section 6 we will give an explicit example by choosing

$$\beta(y) = e^{-ky} \quad (9)$$

Then

$$V(y) = \frac{9}{4} k^2 = \text{const}$$

Equation (5) is simplified further for a particular choice

$$U(y) = \frac{2}{3k} \delta(y) + \epsilon^2 \quad (10)$$

(the reason for the coefficient $2/(3k)$ here will become clear later). With this choice, the field equation and boundary condition read

$$-\ddot{\chi} + \frac{\ddot{a}}{a} \chi - \epsilon^2 p^2 \chi - a^2(\eta) \left[-\chi'' + \frac{9}{4} k^2 \chi \right] = 0 \quad (11)$$

$$\left(\chi' + \frac{3}{2} k \chi - \frac{1}{3k} \frac{p^2}{a^2(\eta)} \chi \right)_{y=0} = 0 \quad (12)$$

Note that even though variables separate in the field equation (11), the dynamics is non-trivial because of the time-dependence of the boundary condition (12).

[§] We are grateful to W. Unruh who pointed out the relevance of this assumption, which simplifies the analysis considerably. We expect, however, that our main results remain valid at $y_0 \gtrsim k^{-1}$ and even $y_0 \gg k^{-1}$. We do not attempt to perform the analysis in the latter cases, as our attitude in this paper is “proof by example”.

3. Static background

In static background, $a(\eta) = 1$, and with $\chi = e^{-i\omega\eta}\chi_\omega$, equation (5) and boundary condition (7) take the form of the static Schrödinger equation

$$-\chi_\omega'' + W_p(y)\chi_\omega = \omega^2\chi_\omega \quad (13)$$

where $\chi(y)$ must be symmetric across the brane, and

$$W_p(y) = V(y) + U(y)p^2$$

(the δ -function term (8) is implicit in $V(y)$). This effective potential has volcano shape, as shown in figure 3. At $p = 0$, continuum of bulk modes starts at $\omega = \frac{3}{2}k$, and there is a bound state with $\omega = 0$ whose wave function is

$$\chi_0(y) = N\beta^{\frac{3}{2}}(y) \quad (14)$$

where N is a normalization factor, ensuring

$$\int_0^\infty |\chi_0(y)|^2 dy = 1$$

This solution corresponds to $\phi_0(y) = \text{const.}$ We assume that this is the only bound state at $p = 0$.




Figure 3. Volcano-shaped effective potential W_p . Its height and large- y asymptotics increase with p , while the dip at $y = 0$ is independent of p .

By continuity, at $p \ll k$ the bound state is still present. Depending on the shapes of $U(y)$ and $V(y)$, there may or may not exist quasi-localized modes at $p \sim k$. For sufficiently small width of the “potential” $U(y)$ (see (6)), at $p \gg k$ even quasi-localized modes cease to exist, while continuum of bulk modes starts at $\omega^2 = \epsilon^2 p^2 + \frac{9}{4}k^2$. These properties are illustrated in Appendix.

As an example, the localized solution to equations (11) and (12) has the form

$$\chi_0 = N \exp \left[- \left(\frac{3}{2}k - \frac{p^2}{3k} \right) y \right]$$

and exists only for

$$p < \frac{3}{\sqrt{2}}k$$

At $p \ll k$ the dispersion relation is $\omega^2 = p^2[1 + O(\epsilon^2)]$ which explains the coefficient in equation (10).

4. Inflation with time-independent parameters

4.1. Preliminaries

At large negative times, the third term in equation (5) dominates, so all modes are shifted in the bulk towards large y , where $U(y) = \epsilon^2$. For given p , all frequencies tend to $\omega = \epsilon p$ as $\eta \rightarrow -\infty$. Thus, at large negative times the evolution proceeds in the bulk and is adiabatic; the field χ has well defined decomposition into positive- and negative-frequency components,

$$\chi = \int \frac{d\lambda}{\sqrt{2\pi\omega_{\lambda,p}}} \left[\chi_{\lambda,p} \exp \left(-i \int \omega_{\lambda,p} d\eta \right) A_{\lambda,p}^- + \text{h.c.} \right] \quad (15)$$

where $\omega_{\lambda,p}^2(\eta)$ and $\chi_{\lambda,p}(y; \eta)$ are eigenvalues and properly normalized eigenstates of the operator

$$U(y)p^2 + a^2(\eta)[- \partial_y^2 + V(y)] \quad (16)$$

and $A_{\lambda,p}^\pm$ are creation and annihilation operators. The asymptotics of $\chi_{\lambda,p}$ at large y is

$$\chi_{\lambda,p} \rightarrow \cos(\lambda y + \varphi_{\lambda,p})$$

where the phase $\varphi_{\lambda,p}$ depends on η . Thus,

$$\omega_{\lambda,p}^2(\eta) = a^2(\eta) \left(\lambda^2 + \frac{9}{4}k^2 \right) + \epsilon^2 p^2 \quad (17)$$

As the physical momenta

$$P(\eta) = \frac{p}{a(\eta)}$$

get redshifted, the properties of modes change. At late times the third term in equation (5) is negligible for superhorizon modes ($P \ll H$), and besides the bulk modes, there is a localized mode

$$\chi_{\text{loc}}(y, \eta) = N \beta^{\frac{3}{2}}(y) \psi_0(\eta) \quad (18)$$

where $\psi_0(\eta)$ obeys the usual equation for superhorizon modes in four dimensions,

$$-\ddot{\psi}_0 + \frac{\ddot{a}}{a} \psi_0 = 0$$

and has a growing component

$$\psi_0 = \phi_0 \cdot a(\eta) \quad (19)$$

where ϕ_0 is independent of time. This is the standard superhorizon mode. It is localized on the brane, the wave function in the transverse dimension being given by equation (14). In terms of the original field ϕ , we have for the localized part

$$\phi_{\text{loc}} = \phi_0$$

Our goal is to relate the constant ϕ_0 in equation (19) to the creation and annihilation operators entering equation (15). This analysis is different for $k \gg H$ and $k \lesssim H$.

4.2. $k \gg H$

It is straightforward to see that in the case $k \gg H$, the evolution of the scalar field is adiabatic when the physical momentum $P(\eta)$ becomes of order k , since at that time the physical frequencies are high,

$$\Omega(\eta) \equiv \frac{\omega(\eta)}{a(\eta)} \gg H$$

and slowly varying,

$$\frac{|\dot{\Omega}|}{a\Omega^2} \lesssim \frac{H}{k}$$

At the time when $P \sim k$, a bound state with $\Omega \sim k \gg H$ emerges in the spectrum of the operator (16), and then its frequency $\omega(\eta)$ slowly decreases from $\omega \sim ak$ at the time when $P \sim k$ to $\omega = p$ at later times when $P \ll k$ but still $P \gg H$. In this time interval the evolution is still adiabatic. Since there is a gap of order k between positive- and negative-frequency parts of the spectrum, there is no mixing between the positive- and negative-frequency modes at the time when the bound state appears, so the bound state is initially in the adiabatic vacuum, up to corrections suppressed by H/k . Therefore, its further evolution proceeds in the same way as in the standard four-dimensional scenario; at the horizon crossing, $P \sim H$, the shape of the bound state is already given by equation (18) modulo H/k -corrections. It is important here that by the time of the horizon crossing, there is no cross talk between the bound state and bulk modes, since they are also separated by the gap of order k (in terms of physical frequencies).

We conclude that for $k \gg H$, the spectrum and amplitude of perturbations coincide with the standard predictions up to corrections suppressed by H/k .

4.3. $k \lesssim H$

In this case the physical momentum crosses H at the same time as it crosses k , or earlier. Adiabaticity gets violated at the same time or before the bound state appears. This case is hard to treat in detail (see, however, sections 5 and 6), but for pure de Sitter

inflation, equation (3), there is a simple scaling argument showing that the spectrum in the localized mode is flat. Let us introduce a new time variable

$$\zeta = p\eta$$

Equation (5) written in terms of ζ does not contain p explicitly. The initial condition is

$$\chi \propto \frac{1}{\sqrt{\epsilon p}} e^{-i\epsilon\zeta} \int d\lambda (\chi_\lambda(y, \zeta) A_{\lambda,p}^- + \text{h.c.})$$

where the factor $(\epsilon p)^{-1/2}$ comes from $(\omega_{\lambda,p})^{-1/2}$ in equation (15), and we recalled that $\omega_{\lambda,p} = \epsilon p$ as $\eta \rightarrow -\infty$. Thus, up to overall factor $p^{-1/2}$, the initial condition does not contain p explicitly either. The boundary condition (7) is independent of p . So, the solution to equation (5) has the general form

$$\chi = \frac{1}{\sqrt{p}} F(\zeta)$$

where for given ζ , the operator F is a linear combination of $A_{\lambda,p}^\pm$ with p -independent coefficients. Therefore, towards the end of inflation, the localized part of the solution, for subhorizon three-momenta, behaves as follows (see equation (18)),

$$\chi_{\text{loc}} = \psi_0(\eta) \chi_0(y)$$

where $\chi_0(y)$ is given by equation (14) and

$$\begin{aligned} \psi_0(\eta) &= \frac{1}{\sqrt{p}} \frac{1}{\zeta} \hat{O}_{\mathbf{p}} \\ &= \frac{1}{p^{3/2}} \frac{1}{\eta} \hat{O}_{\mathbf{p}} \end{aligned}$$

where $\hat{O}_{\mathbf{p}}$ is a time-independent linear combination of $A_{\lambda,p}^\pm$ with p -independent coefficients. The latter property implies that $\hat{O}_{\mathbf{p}}$ describes a Gaussian field obeying

$$\langle \hat{O}_{\mathbf{p}} \hat{O}_{\mathbf{p}'} \rangle = C \delta^3(\mathbf{p} - \mathbf{p}')$$

where C is independent of p . Recalling the relation (4), we obtain that the localized modes

$$\phi_0(\mathbf{p}) \equiv \psi_0(\mathbf{p}) a^{-1}(\eta)$$

towards the end of inflation make up a Gaussian field with flat spectrum,

$$\langle \phi_0(\mathbf{p}) \phi_0(\mathbf{p}') \rangle = \frac{\text{const}}{p^3} \delta^3(\mathbf{p} - \mathbf{p}') \quad (20)$$

It is clear that this relation holds only for momenta which are superhorizon by the end of inflation, higher momenta being suppressed in the usual way. Therefore, if the overall amplitude of perturbations is much smaller than 1, their back reaction on the background geometry is likely to be small. The latter remark applies also to more interesting cases studied in section 5.

It is also clear that for $k \lesssim H$, the constant in equation (20) depends on k and H in a non-trivial way (see also sections 5 and 6). The latter observation implies, in particular, that for non-de Sitter inflation, the expression for the tilt does not generally coincide with that in the standard scenario.

5. Inflation with slowly varying k

5.1. Decaying k

Let us now consider inflation with time-dependent bulk parameters. As an example, let us assume that k slowly decreases, while, for definiteness, H stays constant at inflation. Let k cross H at some $\eta = \eta_\times$, as shown in figure 4. We assume in what follows that k is not grossly different from H after $\eta_\times$, and will not count factors of H/k . On the other hand, we consider ϵ as a small parameter, and will be interested in the dependence of perturbations on this parameter. The analysis of modes with low and high three-momenta gives different results.




Figure 4. Decaying k .

(i) **Low three-momenta.** These are modes that cross out the de Sitter horizon before the time $\eta_\times$, i.e.,

$$P(\eta_\times) \equiv \frac{p}{a(\eta_\times)} \ll H \quad (21)$$

The analysis of these modes goes through as in section 4.2, the spectrum is flat and the amplitude of perturbations in the localized mode is given by the standard four-dimensional formula

$$\phi_0(p) \sim \frac{H}{p^{3/2}} \quad (22)$$

(ii) **High three-momenta.** These modes obey

$$\epsilon P(\eta_\times) \gg H \quad (23)$$

We note in passing that there is an intermediate range of three-momenta, such that $H \ll P(\eta_\times) \ll H\epsilon^{-1}$. This range is actually wide, since we assume $\epsilon \ll 1$.

High-momentum modes (23) are adiabatic at $\eta = \eta_\times$. We will see that the interesting range of transverse momenta is $\lambda \ll k$. Then the interesting modes get out of the adiabatic regime at the time η_ϵ when

$$\epsilon P(\eta_\epsilon) \sim H \quad (24)$$

(we call this moment “ ϵ -horizon crossing”; note that this moment occurs after k crosses H , i.e., $\eta_\epsilon > \eta_\times$). At this time $P(\eta_\epsilon) \gg k$, so for $\lambda \lesssim k$ the modes are still far away from the brane, and the frequencies are all of order ϵp . Just before the ϵ -horizon crossing, the frequencies (17) are

$$\omega_{\lambda,p} = \epsilon p \sqrt{1 + \frac{\lambda^2 + \frac{9}{4}k^2}{H^2}} \quad (25)$$

where we made use of equation (24) to write

$$a(\eta_\epsilon) = \frac{\epsilon p}{H}$$

In what follows we will often omit the second factor in equation (25), since it is of order 1.

The frequency (25) is of order $a(\eta_\epsilon)H$, which means that adiabaticity indeed just becomes violated. According to equation (15), the amplitude of χ at this moment for all interesting λ is

$$\begin{aligned} \chi(\eta_\epsilon) &\propto \frac{1}{\sqrt{\omega}} \\ &\propto \frac{1}{(\epsilon p)^{1/2}} \end{aligned}$$

During some time after ϵ -horizon crossing, the modes are still away from the brane, so equation (5) may be approximated as (modulo mixing^{||} between modes with different λ)

$$-\ddot{\chi}_\lambda + \frac{2}{\eta^2} \chi_\lambda - \frac{1}{\eta^2} \frac{\lambda^2 + \frac{9}{4}k^2}{H^2} \chi_\lambda = 0$$

For $\lambda \ll k$ (and $k < H$, recall that $\eta_\epsilon > \eta_\times$), this equation has both decaying and growing solutions. The growing solution, which is of primary interest, behaves as follows,

$$\chi_\lambda \propto \eta^{-\alpha(\lambda)+1/2} \quad (26)$$

with

$$\alpha(\lambda) = \sqrt{\frac{9}{4} - \frac{\lambda^2 + \frac{9}{4}k^2}{H^2}}$$

This behaviour terminates at the time η_k when

$$P(\eta_k) \equiv \frac{p}{a(\eta_k)} \sim k \quad (27)$$

^{||} This mixing can be shown not to change the estimates that follow, provided that k is not grossly different from H , which we assume throughout.

Around this time the wave functions change considerably; in particular, the localized mode emerges.

The amplitude of χ at this moment is

$$\begin{aligned}\chi_\lambda(\eta_k) &\sim \chi(\eta_\epsilon) \left(\frac{\eta_\epsilon}{\eta_k} \right)^{\alpha(\lambda)-1/2} \\ &\sim \frac{1}{(\epsilon p)^{1/2}} \left(\frac{H}{\epsilon k} \right)^{\alpha(\lambda)-1/2}\end{aligned}$$

With $\epsilon \ll 1$, the integral over the transverse momenta λ is saturated at $\lambda \ll k$, which justifies the assumption made earlier. Since all relevant parameters (P , k and H) are roughly of the same order of magnitude at this time, the estimate of the amplitude of the localized mode at $\eta \sim \eta_k$ is, up to logarithms,

$$\chi_{\text{loc}}(\eta_k) \sim \frac{1}{\sqrt{p}} \frac{1}{\epsilon^\alpha} f\left(\frac{H}{k}\right) \cdot \chi_0(y)$$

where $\chi_0(y)$ is still given by equation (14),

$$\alpha \equiv \alpha(\lambda = 0) = \frac{3}{2} \sqrt{1 - \frac{k^2}{H^2}}$$

and $f(H/k)$ is roughly of order 1. Let us note that for very small three-momenta, the value of k entering here is the asymptotic value at $\eta \gg \eta_\times$.

Later on, at $\eta > \eta_k$, the localized mode grows as $a(\eta)$ (since $k < H$, this mode is superhorizon at η_k and later). Hence, towards the end of inflation the amplitude of the localized mode is (we use the notation (18))

$$\psi_0(\eta) \sim \frac{1}{\sqrt{p}} \frac{1}{\epsilon^\alpha} \frac{a(\eta)}{a(\eta_k)} f\left(\frac{H}{k}\right)$$

Recalling equation (27), we obtain in terms of the original field ϕ that the localized mode of perturbations stays frozen, and its amplitude is

$$\phi_0(p) = \frac{H}{p^{3/2}} \frac{1}{\epsilon^\alpha} \tilde{f}\left(\frac{H}{k}\right) \quad (28)$$

where $\tilde{f}$ is again roughly of order 1. We obtain that at high momenta obeying the relation (23), the spectrum is again flat, while the amplitude is enhanced by the factor

$$\epsilon^{-\alpha} \quad (29)$$

as compared to the modes in the low momentum region, equation (22). Note that $0 < \alpha \leq 3/2$ with $\alpha \rightarrow 3/2$ as $k/H \rightarrow 0$.

The overall behaviour of perturbations is shown in figure 1. Let us make several remarks concerning this result. First, the enhancement of the amplitude, as given by (29), is a free parameter, and can be of several orders of magnitude. The reason for this enhancement is clear: modes with low three-momentum become “tachyonic” when they are still living in the bulk. For $\epsilon = 0$ and constant $k < H$, all modes are tachyonic “from the beginning”, the situation reminiscent of that in four-dimensional theory with the Corley–Jacobson dispersion relation [1, 2]. With our setup, we effectively cut off




Figure 5. Increasing k .

the divergence associated with the tachyonic instability. Second, the beginning of the transition region $p_{\times}$ is determined by the time $\eta_{\times}$ when k crosses H . Clearly, the physically interesting situation occurs when this time is not very distant from the time inflation ends. Third, the momenta in the end and in the beginning of the transition region differ at least by a factor ϵ^{-1} (this is the case shown in figure 1), but their ratio may be much larger, since it is not necessarily determined by the parameter ϵ only, as one might guess by comparing the relations (21) and (23). The value of the parameter k entering the exponent α is to be evaluated somewhere in the interval $(\eta_{\epsilon}(p), \eta_k(p))$, so for very slowly varying $k(\eta)$ this value in fact depends on three-momentum p and settles down only as $k(\eta_k(p))$ reaches its asymptotic value. Fourth, the parameter ϵ may itself depend on time, which would lead to even more complicated spectrum of perturbations. Finally, although we have considered an example with k crossing H , an interesting spectrum is obtained also if $k < H$ during entire inflationary epoch, provided that k and/or ϵ depend on time. The latter two points are clear from equation (28), which is valid for $k < H$, and the above observation that ϵ and k entering equation (28) depend on three momenta, as they are to be evaluated at some η belonging to the interval $(\eta_{\epsilon}(p), \eta_k(p))$.

5.2. Increasing k

If k increases in time as shown in figure 5, the properties of soft and hard modes (in three-dimensional sense) are reversed. **Soft modes** in this case are those for which the bound state gets formed before k crosses H , i.e.

$$P(\eta_{\times}) \equiv \frac{p}{a(\eta_{\times})} \ll k$$

It is these modes that become tachyonic when still living in the bulk. They increase as given by equation (26) from the time η_{ϵ} to η_k ; these times are still determined by

equations (24) and (27), with k now being the value of $k(\eta)$ in the past, where $k < H$. At the time η_k the bound state emerges, and then the amplitude of the bound state mode increases as $a(\eta)^{-1}$. Thus, equation (28) applies now to the soft modes.

Hard modes are those which never become tachyonic when staying in the bulk, which is the case for

$$\epsilon P(\eta_x) \gg H$$

They start growing after forming the bound state, so their amplitudes are given by the four-dimensional expression (22). The overall spectrum is thus the reverse of figure 1; it is shown in figure 2. The remarks made in the end of section 5.1 apply to the case of increasing k as well.

6. Explicit example

In this section we consider an explicit setup outlined in the end of section 2. The field χ obeys equation (11) with the boundary condition (12). We will be interested in the case $k \sim H$ and, as in section 5, study the dependence of the amplitudes on ϵ , assuming $\epsilon \ll 1$. As mentioned in section 2, in spite of the fact that variables separate in equation (11), the dynamics is non-trivial due to the time-dependence of the boundary condition (12). We simplify the analysis further by making use of the instantaneous approximation to the boundary condition (12). Namely, at $\eta < \eta_k$, where η_k is given by equation (27), i.e.

$$\eta_k = -\frac{k}{Hp} \sim -\frac{1}{p} \quad (30)$$

the third term in equation (12) dominates and we omit the first two terms. So, at these times the boundary condition reads

$$\chi|_{y=0} = 0 \quad \text{at } \eta < \eta_k \quad (31)$$

On the other hand, at $\eta > \eta_k$ we neglect the third term in equation (12) and the boundary condition becomes

$$\left(\chi' + \frac{3}{2}k\chi \right)_{y=0} = 0 \quad \text{at } \eta > \eta_k \quad (32)$$

In this approximation, the problem reduces to that of solving the field equation (11) with the boundary conditions (31), (32) and matching the solutions at $\eta = \eta_k$.

At $\eta < \eta_k$ the solution is (we slightly change the notation as compared to equation (15))

$$\chi = \int_0^\infty \frac{d\lambda}{\sqrt{2\pi}} [\chi_{\lambda,p}(y, \eta) A_{\lambda,p}^- + \text{h.c.}]$$

where $\chi_{\lambda,p}$ are

$$\chi_{\lambda,p}(y, \chi) = -i\sqrt{\pi} e^{\frac{i\pi\alpha}{2}} \sqrt{-\eta} H_\alpha^{(1)}(-\eta\epsilon p) \sin(\lambda y) \quad (33)$$

and

$$\alpha = \sqrt{\frac{9}{4} \left(1 - \frac{k^2}{H^2}\right) - \frac{\lambda^2}{H^2}} \quad (34)$$

As $\eta \rightarrow -\infty$, these modes approach properly normalized negative-frequency plane waves

$$\chi_{\lambda,p}(y, \eta) = \sqrt{\frac{2}{\omega}} e^{-i\omega\eta + i\varphi} \sin(\lambda y)$$

where $\omega = \epsilon p$ and φ is a constant phase. For $k < H$ and small enough λ the parameter α is real. So, these modes indeed grow like $|\eta|^{1/2-\alpha}$ in the interval $\eta_\epsilon \lesssim \eta \lesssim \eta_k$, where η_ϵ is determined by equation (24). For $k > H$, the parameter α is imaginary, so the modes (33) oscillate in η for all $\eta < \eta_k$.

At late times, $\eta > \eta_k$, besides the bulk modes, there are two localized modes obeying equations (11) and (32),

$$\chi_{\text{loc}}^\pm(y, \eta) = \sqrt{3k} \cdot \sqrt{-\eta} (-\eta)^{\pm \frac{3}{2}} e^{-\frac{3}{2}ky}$$

These are the standard four-dimensional modes; recall that $k \sim H$, so at $\eta > \eta_k$ these modes are superhorizon. At $\eta > \eta_k$ the field χ is a superposition

$$\chi(y, \eta) = C^{(-)} \chi_{\text{loc}}^-(y, \eta) + C^{(+)} \chi_{\text{loc}}^+(y, \eta) + \text{bulk modes} \quad (35)$$

The coefficients here are obtained by matching this solution to the solution (33) at $\eta = \eta_k$. We are interested in the coefficient $C^{(-)}$, which is the amplitude of the growing localized mode. To this end we notice that the bulk modes entering equation (35) are orthogonal to $\chi_{\text{loc}}^\pm$ with the standard scalar product

$$(\psi, \chi) = \int_0^\infty dy (\dot{\chi} \psi^* - \dot{\psi}^* \chi)$$

On the other hand,

$$(\chi_{\text{loc}}^+, \chi_{\text{loc}}^-) = 3$$

Thus

$$C^{(-)} = \frac{1}{3} (\chi_{\text{loc}}^+, \chi)_{\eta=\eta_k}$$

where χ is the solution (33), and the scalar product is to be evaluated at $\eta = \eta_k$. One obtains for arbitrary η

$$(\chi_{\text{loc}}^+, \chi) = \sqrt{3k\pi} \frac{(-\eta p \epsilon)^{\frac{3}{2}} e^{\frac{i\pi\alpha}{2}}}{(p\epsilon)^{\frac{3}{2}}} \left(\frac{3}{2} H_\alpha^{(1)}(-\eta\epsilon p) - \eta \dot{H}_\alpha^{(1)}(-\eta\epsilon p) \right) \frac{i\lambda}{\lambda^2 + \frac{9}{4}k^2}$$

Thus

$$C^{(-)} = \frac{1}{\sqrt{3kp}^{\frac{3}{2}}} \int_0^\infty \frac{d\lambda}{\sqrt{2}} e^{\frac{i\pi\alpha}{2}} \left(\frac{3}{2} H_\alpha^{(1)}(\epsilon) - \epsilon \partial_\epsilon H_\alpha^{(1)}(\epsilon) \right) \frac{ik\lambda}{\lambda^2 + \frac{9}{4}k^2} A_{\lambda,p}^- + \text{h.c.} \quad (36)$$

where we took into account equation (30).

Now we are ready to calculate the spectrum of perturbations in the localized mode, defined as

$$\langle \phi_0(\mathbf{p}) \phi_0(\mathbf{p}') \rangle \equiv \frac{2\pi^2}{p^3} \mathcal{P}(p) \delta^3(\mathbf{p} - \mathbf{p}')$$

We make use of the relation (4) of the original field ϕ to the field χ and the standard commutation relations

$$[A_{\lambda,p}^-, A_{\lambda',p'}^+] = \delta^3(\mathbf{p} - \mathbf{p}') \delta(\lambda - \lambda')$$

With the expression (36) for the coefficient $C^{(-)}$ we find

$$\mathcal{P}(p) = \frac{H^2}{4\pi^2} \int_0^\infty d\lambda e^{\frac{i\pi(\alpha-\alpha^*)}{2}} \frac{\lambda^2 k}{3(\lambda^2 + \frac{9}{4}k^2)^2} \left| \frac{3}{2} H_\alpha^{(1)}(\epsilon) - \epsilon \partial_\epsilon H_\alpha^{(1)}(\epsilon) \right|^2 \quad (37)$$

For time-independent k , the right hand side of equation (37) is independent of p , the spectrum is flat, while the amplitude depends on k , H and ϵ in a non-trivial way. This is in accord with the general analysis given in section 4.

On the other hand, if k slowly varies in time, the spectrum is no longer flat. The important time interval is $\eta_\epsilon < \eta < \eta_k$, in which bulk modes grow provided that $k < H$, and do not grow for $k > H$.

Let us consider the case of decreasing k , figure 4. For low three-momenta one has (see equation (21)) $\eta_\times \gg \eta_k$ and $\alpha(k)$ defined in (34) is pure imaginary. So, the integrand in equation (37) is independent of ϵ at small ϵ . The spectrum is flat, and the power is

$$\mathcal{P}(p) = \frac{H^2}{4\pi^2} \cdot \text{const},$$

where the constant is of order 1. In the opposite case of high three-momenta, the times are ordered as $\eta_\times < \eta_\epsilon < \eta_k$. So, $k(\eta_k) < H$ and $\alpha(k, \lambda)$ is real at small enough λ . In this case

$$\left| \frac{3}{2} H_\alpha^{(1)}(\epsilon) - \epsilon \partial_\epsilon H_\alpha^{(1)}(\epsilon) \right|^2 = \frac{1}{4\pi} \left(\frac{2}{\epsilon} \right)^{2\alpha} \left(\alpha + \frac{3}{2} \right)^2 \frac{\Gamma(\alpha)^2}{\Gamma(3/2)^2}$$

Substituting this expression into equation (37) one finds that the integral is saturated at small transverse momenta, $\lambda^2 \sim H^2 / \ln \epsilon$. Therefore, we again obtain the flat spectrum but with the amplitude enhanced by the factor (29) (up to logarithm), in agreement with the discussion in section 5.1.

If k increases in time as shown in figure 5, $\alpha(k)$ is imaginary for high momenta ($\eta_\times > \eta_k$) and real for low momenta ($\eta_\times < \eta_\epsilon$). The perturbations get enhanced by the factor $\epsilon^{-\alpha}$ at low momenta, and do not get enhanced at high momenta, so we obtain the spectrum shown in figure 2.

Acknowledgments

The authors are indebted to S. Dubovsky, M. Sazhin, M. Shaposhnikov, S. Sibiryakov and W. Unruh for helpful discussions and comments. This work is supported in part

by RFBR grant 05-02-17363-a, by the Grants of the President of Russian Federation NS-2184.2003.2, MK-3507.2004.2, by INTAS grant YSF 04-83-3015, and by the grant of the Russian Science Support Foundation.

Appendix

In this Appendix we illustrate the properties of eigenmodes of equation (13) by considering a step-function “potential”

$$U(y) = \alpha^2 \theta(y_0 - |y|) + \epsilon^2 \theta(|y| - y_0)$$

where $\alpha \sim 1$ and $\epsilon \ll 1$. We take the warp factor $\beta(y)$ in the form (9), so that another “potential” is constant,

$$V(y) = \frac{9}{4}k^2$$

We consider equation (13) on half-line $y > 0$ and impose the boundary condition (7), i.e.,

$$\left(\chi' + \frac{3}{2}k\chi \right)_{y=0} = 0 \quad (38)$$

We are interested in localized or quasilocalized modes whose general expression at $y < y_0$ is

$$\chi_\omega = Ae^{-\nu y} + Be^{\nu y}, \quad y < y_0 \quad (39)$$

Then

$$\omega^2 = -\nu^2 + \frac{9}{4}k^2 + \alpha^2 p^2 \quad (40)$$

while the boundary condition (38) reads

$$\left(\frac{3}{2}k - \nu \right) A + \left(\frac{3}{2}k + \nu \right) B = 0 \quad (41)$$

Let us begin with localized modes. They decay at $y > y_0$,

$$\chi_\omega = e^{-\lambda(y-y_0)}, \quad y > y_0$$

where we set the overall factor equal to one. From equations (13) and (40) we obtain

$$\nu^2 = \lambda^2 + \hat{\alpha}^2 p^2 \quad (42)$$

where $\hat{\alpha}^2 = \alpha^2 - \epsilon^2$. By matching χ_ω and χ'_ω at $y = y_0$ we find

$$A = \frac{\nu + \lambda}{2\nu} e^{\nu y_0}$$

$$B = \frac{\nu - \lambda}{2\nu} e^{-\nu y_0}$$

and the relation (41) gives

$$e^{-2\nu y_0} = \frac{\nu + \lambda}{\nu - \lambda} \cdot \frac{\nu - \frac{3}{2}k}{\nu + \frac{3}{2}k} \quad (43)$$

This equation, supplemented with equation (42), determines the eigenvalue ν for given three-momentum p .

At $p^2 = 0$ equation (13) has a zero mode $\chi_0 = \exp(-\frac{3}{2}ky)$ with $\lambda = \nu = \frac{3}{2}k$ and $\omega = 0$. For $p^2 \ll k^2$ there exists a solution to equations (42) and (43) describing a localized state close to the zero mode,

$$\begin{aligned}\nu &= \frac{3}{2}k + \frac{\hat{\alpha}^2}{3k}e^{-3ky_0} \cdot p^2 + O(p^4) \\ \omega^2 &= (\alpha^2 - \hat{\alpha}^2 e^{-3ky_0}) \cdot p^2 + O(p^4)\end{aligned}$$

Up to rescaling of coordinates on the brane, this mode has relativistic dispersion relation.

At $p^2 \gg k^2$ no real solutions to equations (42), (43) exist: equation (42) gives $\nu \gtrsim p$, so the right hand side of equation (43) is greater than 1 in that case. Thus, there are no localized modes at high three-momenta p .

Let us now turn to quasilocalized modes. They still have the form (39) at $y < y_0$, while at $y > 0$ they oscillate

$$\chi_\omega = be^{iq(y-y_0)} + b^*e^{-iq(y-y_0)}$$

so that

$$q^2 = \hat{\alpha}^2 p^2 - \nu^2$$

The normalization condition appropriate for the modes in continuum is

$$|b|^2 = \frac{1}{2} \tag{44}$$

By matching χ_ω and χ'_ω at $y = y_0$ we obtain

$$b = \frac{1}{2} (Ae^{-\nu y_0} + Be^{\nu y_0}) + \frac{i\nu}{2q} (Ae^{-\nu y_0} - Be^{\nu y_0})$$

so the normalization condition (44) and equation (41) determine the coefficient A ,

$$|A|^2 = 2 \left[\left(e^{-\nu y_0} + \frac{\nu - \frac{3}{2}k}{\nu + \frac{3}{2}k} e^{\nu y_0} \right)^2 + \frac{\nu^2}{q^2} \left(e^{-\nu y_0} - \frac{\nu - \frac{3}{2}k}{\nu + \frac{3}{2}k} e^{\nu y_0} \right)^2 \right]^{-1} \tag{45}$$

The defining property of a quasilocalized mode is that it is large at the brane position (when normalized according to (44)),

$$|A| \gg 1 \tag{46}$$

At $ky_0 \gg 1$ this relation indeed holds for

$$\nu = \frac{3}{2}k + O(e^{-3ky_0})$$

In that case the quasilocalized mode exists for arbitrarily high three-momenta p ; furthermore, at $p \gg k$ and

$$\nu = \frac{3}{2}k - 3ke^{-3ky_0} + O\left(\frac{k^2}{p}e^{-3ky_0}\right)$$

one has

$$|A| \sim \frac{p}{k} e^{3ky_0/2}$$

so the localization gets stronger at high momenta p .

At $ky_0 \lesssim 1$ the only way the relation (46) may be satisfied is that p is large, and the first term in the square bracket in (45) vanishes, which gives

$$e^{-2\nu y_0} = \frac{\frac{3}{2}k - \nu}{\frac{3}{2}k + \nu}$$

The latter equation, however, has positive solutions only for

$$y_0 > \frac{2}{3k}$$

Thus, a quasilocalized mode exists at high three-momenta p provided that ky_0 is large enough, otherwise there are no quasilocalized modes at $p \gg k$. It is the latter case that is considered in the main text.

References

- [1] Martin J and Brandenberger R H 2001 *Phys. Rev. D* **63** 123501
- [2] Martin J and Brandenberger R H 2001 *Mod. Phys. Lett. A* **16** 999
- [3] Niemeyer J C 2001 *Phys. Rev. D* **63** 123502
- [4] Starobinsky A A 2001 *Pisma Zh. Eksp. Teor. Fiz.* **73** 415 [*JETP Lett.* **73** 371]
- [5] Lemoine M, Lubo M, Martin J and Uzan J P 2002 *Phys. Rev. D* **65** 023510
- [6] Brandenberger R H, Joras S E and Martin J 2002 *Phys. Rev. D* **66** 083514
- [7] Martin J and Brandenberger R H 2002 *Phys. Rev. D* **65** 103514
- [8] Danielsson U H 2002 *Phys. Rev. D* **66** 023511
- [9] Danielsson U H 2002 *J. High Energy Phys.* JHEP07(2002)040
- [10] Bozza V, Giovannini M and Veneziano G 2003 *Journal of Cosmology and Astroparticle Physics* JCAP05(2003)001
- [11] Martin J and Brandenberger R 2003 *Phys. Rev. D* **68** 063513
- [12] Kempf A 2001 *Phys. Rev. D* **63** 083514
- [13] Kempf A and Niemeyer J C 2001 *Phys. Rev. D* **64** 103501
- [14] Easther R, Greene B R, Kinney W H and Shiu G 2001 *Phys. Rev. D* **64** 103502
- [15] Easther R, Greene B R, Kinney W H and Shiu G 2003 *Phys. Rev. D* **67** 063508
- [16] Brandenberger R and Ho P M 2002 *Phys. Rev. D* **66** 023517
- [17] Kaloper N, Kleban M, Lawrence A E and Shenker S 2002 *Phys. Rev. D* **66** 123510
- [18] Schalm K, Shiu G and van der Schaar J P 2004 *J. High Energy Phys.* JHEP04(2004)076
- [19] Lizzi F, Mangano G, Miele G and Peloso M 2002 *J. High Energy Phys.* JHEP06(2002)049
- [20] Tanaka T 2000 A comment on trans-Planckian physics in inflationary universe *Preprint* astro-ph/0012431
- [21] Porrati M 2004 *Phys. Lett. B* **596** 306
- [22] Brandenberger R H and Martin J 2005 *Phys. Rev. D* **71** 023504
- [23] Corley S and Jacobson T 1996 *Phys. Rev. D* **54** 1568
- [24] Visser M 1985 *Phys. Lett. B* **159** 22
- [25] Chung D J H and Freese K 2000 *Phys. Rev. D* **61** 023511
- Chung D J H and Freese K 2000 *Phys. Rev. D* **62** 063513
- [26] Chung D J H, Kolb E W and Riotto A 2002 *Phys. Rev. D* **65** 083516
- [27] Csaki C, Erlich J and Grojean C 2001 *Nucl. Phys. B* **604** 312
- [28] Dubovsky S L 2002 *J. High Energy Phys.* JHEP01(2002)012
- [29] Kraus P 1999 *J. High Energy Phys.* JHEP12(1999)011
- [30] Bowcock P, Charmousis C and Gregory R 2000 *Class. Quant. Grav.* **17** 4745
- [31] Sibiryakov S and Gorbunov D 2005 Private communication
- [32] Langlois D, Maartens R and Wands D 2000 *Phys. Lett. B* **489** 259